

## Advanced search

Showing results 101 to 120 on 7065 in total

### EVENTS

Relevance:  100%

## Approche dynamique pour l'interprétation causale dans les études portant sur le vieillissement

We consider the issue of estimating causal effects in a dynamic approach based on a multivariate stochastic process representation, which may be called the "stochastic system approach". Conditional and marginal effects can be defined. We focus on the issue of the horizon on which causal influence must be studied, in particular in ageing studies. In ageing studies, one of the most important events that we have to consider is "death". This is why the illness-death model is important in such studies. But "death" is not an event which is on the same footing as other events that can happen to subjects. Even if the vital status is part of the state, it has a very special meaning, in that all the other components of the state are defined only for a living subject. The consequence is that causal influences must be defined on a maximum horizon which is the time of death. We do not say that death has an influence on the other components of the state, but that these other components are not defined after death. For instance if we are interested in dementia, the state can be represented by a bivariate counting process counting dementia and death. However dementia is defined only for a living subject: after death the subject does not exist anymore and cannot be qualified as demented or not demented. When we investigate the causal influence of a factor, we should first look at its causal influence on death, then on its influences on other processes. Cases where a value of a modifiable factor can be preferred to another one will be given. Thus, the stochastic system approach helps clarifying the important issue of assessing causal effects in ageing studies Contact : Mme Mariethé CHAUMEIL Inscription gratuite mais obligatoire (commande sandwiches avant le lundi 16 MARS 2015 Courriel : mariethe.chaumeil@chu-lyon.fr

06/11/2020

### EVENTS

Relevance:  100%

## Vibrio diversity and oyster mortality: the hypothesis of a new polymicrobial disease

Vibrios have been associated with successive mortality outbreaks of *Crassostrea gigas* in France that have resulted in losses up to 100% of production. Given the near monoculture of *C. gigas* in Europe, there is an urgent need to understand the epidemiology of these outbreaks, particularly the role of *Vibrio* in the diseases. The study of the *Vibrios* distribution on fine phylogenetic and spatial scales has demonstrated that vibrios coexisting in the water column can be divided into closely related populations, which pursue different lifestyles i.e. ecological population (Hunt et al., 2008). However, a link between ecological populations and pathogenicity has not been demonstrated, and it is unclear whether pathogenicity is a trait primarily linked to clones or to populations comprising a large number of distinct genotypes. In the present *Vibrio* populations in an intensive oyster cultivation area. We demonstrate that *Vibrio* populations do not assemble neutrally in oysters from water column populations i.e. specific genotypes colonize the oysters. Combining experimental ecology, high throughput infection assay and genome sequencing, we showed that the onset of disease in oysters is associated with progressive replacement of diverse, benign colonizers by members of a phylogenetically coherent virulent population together with quorum sensing pheromone producers. Analyses of oyster mortality following experimental infection suggest that disease onset can be facilitated by the presence of non-virulent strains. Oyster disease may thus represent a new form of polymicrobial disease, in which non-pathogenic strains contribute to increased mortality. Hunt DE, et al. (2008) Resource partitioning and sympatric differentiation among closely related bacterioplankton. Science 320(5879):1081-1085. Lemire A, Goudenège D, Versigny T, Petton B, Calteau A, Labreuche Y, Le Roux F. (2014) Populations, not clones, are the unit of vibrio pathogenesis in naturally infected oysters. ISME J. Dec 9. doi: 10.1038/ismej.2014.233

06/11/2020

#### EVENTS

Relevance:  100%

## Lessons from 30 years of natural disturbances in the Bavarian Forest National Park

-

06/11/2020

---

#### EVENTS

Relevance:  100%

## Transposable element evolution in Drosophila

-

06/11/2020

---

#### EVENTS

Relevance:  100%

## Similarities and differences between evolutionary processes in linguistics and biology

Among biologist as well as linguists, it is now widely accepted that there are many striking parallels between the evolution of life forms and the history of languages. Starting from the rise of language studies as a scientific discipline in the early 19th century, up to today's recent "quantitative turn" in historical linguistics, scholars from both disciplines have repeatedly pointed to similarities between the respective research objects in biology and linguistics. During the last two decades, this has lead to a new school of "quantitative historical linguistics". Based on the key assumption that the characteristic processes of language change and biological evolution are so similar that the methods designed for one discipline may also be used in the other one, methods which were originally designed to study biological evolution (methods for phylogenetic reconstruction, sequence alignment, or biological network analysis) have now repeatedly been applied to linguistic data. Unfortunately, not all analogies which have been made between evolutionary processes in linguistics and biology reflect true similarities in the processes. Striking differences between the research objects of both disciplines are often ignored. In the talk, I will review proposed similarities between evolutionary processes in the two disciplines and discuss their methodological implications.

06/11/2020

---

#### EVENTS

Relevance:  100%

## Bad luck of cancer - or misinterpreted statistics?

A recent paper in Science (January 2015) claimed that the majority, 64% to be precise, of cancers is due to bad luck, so non-preventable. The message was spread quickly through media, including serious ones like BBC (and also radio Slovenia, if I may add). And while the paper has been criticized by many, the authors seem to stick to the original message. In this talk I'll give my view of the paper and try to defend a counter message that their analysis gives them absolutely no grounds to make such a claim. The arguments that I'll present have, to my knowledge, not appeared in published reactions to the paper. Obviously, if I am right, they are wrong, and vice versa. If I was Bayesian, I would, at present, give a 0.99 prior probability to the first option. Either way, it should be fun.

06/11/2020

---

#### EVENTS

Relevance:  100%

## Human sex differences in height : A costly evolution due to gender hierarchy.

-

06/11/2020

---

#### EVENTS

Relevance:  100%

## Titre à venir

-

06/11/2020

---

#### EVENTS

Relevance:  100%

## Understanding molecular evolution: insights from animals and plants

-

06/11/2020

---

#### EVENTS

Relevance:  100%

## Dispersion des vers de terre en Agroécosystèmes : résultats du projet Edisp

La dispersion est un processus central pour le maintien de la biodiversité dans les paysages. Cependant les patterns de dispersion et leurs déterminants sont encore mal documentés, en particulier chez les organismes du sol. En conséquence, une grande partie des déterminants de la biodiversité des sols n'est toujours pas comprise. Nous exposons les résultats d'un projet ANR de quatre ans sur la dispersion des vers de terre, un organisme clé dans le maintien de nombreux services écosystémiques. Avec une combinaison d'approches (1) de terrain multi-échelle, (2) de (méta) communauté, (3) de génétique du paysage, et (4) d'expérience en laboratoire, nous avons identifié des mécanismes centraux qui permettent aux vers de terre de disperser dans un agro-écosystème de manière adaptée, et nous montrons l'importance de la dispersion pour le maintien de la diversité spécifique et génétique. Nos résultats suggèrent également que la dispersion passive joue un rôle sous estimé dans la dynamique spatiale de ces organismes.

06/11/2020

---

#### EVENTS

Relevance:  100%

## Baiting for phylogenetic fishing

-

06/11/2020

---

#### EVENTS

Relevance:  100%

## Phylogenetic Fate Mapping

We are exploring techniques using single cell genomic sequencing in human beings for the purpose of defining phylogenetic trees to explain the relationship of cells within defined human tissues. I will discuss how we used single cell exome and whole genome sequencing to identify patient and donor adipocytes in human white adipose tissue biopsies taken from bone marrow recipients and discuss future projects using similar methods to explore cellular identities in human tissue samples.

06/11/2020

---

#### EVENTS

Relevance:  100%

## Beyond Jarman-Bell: how principles of digestive physiology can and cannot inform on ecological diversification.

-

06/11/2020

---

#### EVENTS

Relevance:  100%

## Evaluation quantitative et qualitative d'un dispositif de surveillance basé sur le principe de la déclaration obligatoire : l'exemple de la surveillance de la brucellose chez les bovins en France

La surveillance de la brucellose chez les bovins, maladie pour laquelle la France est indemne depuis 2005, a pour objectif principal d'assurer la détection de tout nouveau foyer qui surviendrait sur le territoire. Cette surveillance repose principalement sur la surveillance événementielle (clinique) : celle-ci impose aux éleveurs et aux vétérinaires la déclaration de tout avortement chez les bovins, suivi du dépistage de la femelle ayant avorté vis-à-vis de la brucellose. Toutefois, en amont de ce travail, il était admis par l'ensemble des acteurs, sans que cela ait été rigoureusement évalué, que cette surveillance souffrait d'une forte sous-déclaration et nécessitait d'être améliorée. Les objectifs de l'évaluation de ce dispositif étaient : 1) de quantifier le niveau de sous-déclaration et estimer l'effet de différents facteurs sur le processus de déclaration ; 2) d'analyser le processus de décision conduisant les éleveurs et les vétérinaires à participer ou non au dispositif. Afin de répondre au premier objectif, le principe des méthodes de capture-recapture unilistes a été retenu. L'ensemble des observations a pu être modélisé à l'aide des méthodes MCMC et de deux modèles multi-réponses, le modèle de Poisson enflé en zéro (ZIP) et le modèle hurdle. La deuxième étude a consisté à mener des entretiens semi-directifs auprès de 12 éleveurs et de leurs 8 vétérinaires.

06/11/2020

---

## EVENTS

Relevance:  100%

# Host phylogeny and diet drive the structure of mammalian microbial communities at different phylogenetic scales

Bacterial communities (microbiota) living in Mammal guts are composed of thousands of bacterial species that are essential for host physiology, immunity and diet. It has been shown that host phylogeny (genetics and immune system of the host) and diet are the two major factors driving the composition of gut microbiota. However, major questions remain: (i) the relative contribution of host phylogeny and diet at short and long time scales is highly debated and the two processes are not well characterized, (ii) host phylogeny may drive the composition through co-evolution with bacterial lineages or through niche selection, with closely-related hosts retaining similar bacteria from the environment. Here, we show that host phylogeny and diet are for the most part independent processes and do not drive the bacterial composition at the same taxonomic scale. Diet determines what lineage is present or not at deep bacterial phylogenetic levels through gain or loss of lineages creating nested communities. Host phylogeny, however, selects the lineages at finer scales through true turnover of lineages, consistent with a more stringent selection of tolerated antigens. Finally, it appears that co-speciation between hosts and bacterial lineages plays a minor role in driving the correlation between community composition and host phylogeny, suggesting that environmental filtering by host genetics is the dominant process at selecting bacterial lineages. Our results shed light on the long-timescale evolutionary dynamic of gut bacterial communities, which are multi-layered phylogenetic structures shaped very differently by host phylogeny and diet.

06/11/2020

---

## EVENTS

Relevance:  100%

# Gathering, updating, and calibrating phylogenies for The Open Tree of Life

The Open Tree of Life project is a collaborative effort to synthesize, share, and update a comprehensive phylogeny of all 2.3 million named species. We have completed a draft synthesis of a single tree from hundreds of phylogenetic estimates using taxonomy as a scaffold. This synthesis is not static but rather will be continually revised as new data become available. This undertaking requires development of both novel infrastructure and analysis tools. I will discuss three components of this project: Phylsystem, an open database and web application for community curation of phylogenies using a git-based datastore, PhyScraper, a pipeline to continually update phylogenetic estimates as new data is generated, and FastDate, an algorithm to rapidly generate maximum a posteriori estimates of time-calibrated trees, even for phylogenies with hundreds or thousands of tips. Together, these developments reduce impediments to accessing, analyzing and reusing the phylogenetic information which is essential to biological research today.

06/11/2020

---

## EVENTS

Relevance:  100%

# Transitions between combined and separate sexes in plants

-

06/11/2020

---

## EVENTS

Relevance:  100%

# Characterisation of the bacterial replicons

The genome of bacteria is classically separated into essential, stable and slow evolving replicons (chromosomes) and accessory, mobile and rapidly evolving replicons (plasmids). This paradigm is being questioned since the discovery of genomic elements that possess both chromosomal and plasmidic features. These Extra-Chromosomal Essential Replicons (ECERs), be they called "megaplasids", "secondary chromosomes" or "chromids", are found in diverse lineages across the bacterial phylogeny and are generally believed to be modified plasmids. However, their true nature and the mechanisms permitting their integration within the stable genome are yet to be formally determined. The relationships between replicons, with reference to their Genetic Information Inheritance Systems (GIIS), were explored under the assumption that the inheritance of ECERs is integrated to the cell cycle and highly constrained in contrast to that of standard plasmids. A global comparative genomics analysis including all available complete bacterial genome sequences, was performed using GIIS functional homologues as parameters and applying several analytical procedures. GIIS proved appropriate in characterizing the level of integration within the stable genome, as well as the origins, of the replicons. The study of ECERs thus provides clues to the genetic mechanisms and evolutionary processes involved in the replicon stabilization into the essential genome and to the continuity of the genomic material.

06/11/2020

---

## EVENTS

Relevance:  100%

# A turtle view of the evolution of genomes and sex determination

-

06/11/2020

---

## EVENTS

Relevance:  100%

# Dynamics of slow cell renewal in humans

Human tissues constantly replace dying cells with newborn cells. The pace at which they are replaced, however, varies by orders of magnitudes between blood cells, which are renewed every day and neurons, for which renewal is non-existent or limited to specific regions of the brain. Between those extreme are many tissues that turnover on a time scale of years, although no direct measurements have been done. We present here a mathematical method to estimate cell turnover in slowly renewing biological systems. Age distribution of DNA can be estimated from the integration of radiocarbon derived from nuclear bomb-testing during the cold war (1955-1963). For slowly renewing tissues, this method provides a better estimate of the average age of the tissue than direct estimates from the bomb-curve. Moreover, death, birth and turnover rates can be estimated. We highlight this method with data from hippocampal neurons and cardiomyocytes.

06/11/2020

---

## RESULTS FILTER

## Par types de contenus

---

- [Basic page](#)  
(97)
- [Department](#)  
(5)
- [Directory of people](#)  
(247)
- [Events](#)  
(1228)
- [News](#)  
(356)
- [Photo/video gallery](#)  
(2)
- [Publications](#)  
(5112)
- [Team](#)  
(18)

## Par rubrique

---

- [Recherche](#)  
(22)
- [Intranet](#)  
(21)
- [Laboratoire](#)  
(10)
- [Research](#)  
(3)
- [Life in the lab](#)  
(1)
- [Recherche Recherche](#)  
(1)
- [Vie au laboratoire](#)  
(1)

## By creation date

---

From  
(jj/mm/aaaa)

To  
(jj/mm/aaaa)

FILTER BY  
DATE

## SORT BY :

- > [Relevance](#)
- > [Title](#)
- > [Type of content](#)
- > [Publication date](#)